

FLUORESCENT LY α EMISSION FROM THE HIGH-REDSHIFT INTERGALACTIC MEDIUM

SEBASTIANO CANTALUPO, CRISTIANO PORCIANI, SIMON J. LILLY AND FRANCESCO MINIATI

Institute for Astronomy, ETH Zürich, CH-8093 Zürich, Switzerland
 cantalupo@phys.ethz.ch, porciani@phys.ethz.ch, lilly@phys.ethz.ch, fm@phys.ethz.ch

Draft version February 2, 2008

ABSTRACT

We combine a high-resolution hydro-simulation of the Λ CDM cosmology with two radiative transfer schemes (for continuum and line radiation) to predict the properties, spectra and spatial distribution of fluorescent Ly α emission at $z \sim 3$. We focus on line radiation produced by recombinations in the dense intergalactic medium ionized by UV photons. In particular, we consider both a uniform background and the case where gas clouds are illuminated by a nearby quasar. We find that the emission from optically thick regions is substantially less than predicted from the widely used static, plane-parallel model. The effects induced by a realistic velocity field and by the complex geometric structure of the emitting regions are discussed in detail. We make predictions for the expected brightness and size distributions of the fluorescent sources. Our results account for recent null detections and can be used to plan new observational campaigns both in the field (to measure the intensity of the diffuse UV background) and in the proximity of bright quasars (to understand the origin of high column-density absorbers).

Subject headings: cosmology: theory – intergalactic medium – large-scale structure of universe – line: formation – quasars: absorption lines – radiative transfer

1. INTRODUCTION

Hydrogen absorption-line systems observed shortward of Ly α emission in quasar spectra constitute an important probe of the physical state of the intergalactic medium at high-redshift. These spectral features are shaped by the combined action of gravity, hydrodynamics and photoionization processes which determine the local density and the velocity field of neutral hydrogen within the absorbers. Numerical simulations suggest that the so called Lyman- α forest is generated by diffuse, sheetlike and filamentary structures with a mean density which is between 1 and 10 times higher than the cosmic average (Cen et al. 1994; Zhang, Anninos & Norman 1995; Hernquist et al. 1996; Miralda-Escudé et al. 1996). These low-column-density systems are highly ionized by the extragalactic background of Lyman continuum photons generated by young stellar populations and quasars. At the opposite extreme, Lyman-limit (LLS, $N_{\text{HI}} > 10^{17.2} \text{ cm}^{-2}$) and damped Lyman- α (DLA, $N_{\text{HI}} > 10^{20.3} \text{ cm}^{-2}$) systems correspond to concentrations of atomic hydrogen which are optically thick to the cosmic ionizing background. Numerical simulations suggest that they arise in dense gas clouds with a meatball topology. On cosmological scales, they appear to form a collection of isolated clouds which trace the cosmic web.

Optically thick clouds are expected to emit fluorescent Ly α photons produced in hydrogen recombinations (Hogan & Weymann 1987; Gould & Weinberg 1996). This emission is concentrated in the outer parts of the clouds where hydrogen is significantly ionized by the external UV background ($\tau_{\text{LL}} \sim 1$). However, Ly α photons cannot directly escape the clouds because of the large optical depth in the center of the line ($\tau_{\text{Ly}\alpha} \simeq 10^4 \tau_{\text{LL}}$). Each photon thus suffers a large number of resonant scatterings (more precisely: absorptions and re-emissions) by neutral hydrogen atoms in the ground state. Each scat-

tering adds a small Doppler shift to the frequencies of the photons due to the thermal (and turbulent) motions of the atoms. Therefore, photons execute a random walk both in frequency and in physical space until their frequencies are shifted sufficiently away from the line center and they are able to escape the medium in a single flight (Zanstra 1949).

Monte Carlo simulation (e.g. Ahn, Lee & Lee 2001; Zheng & Miralda-Escudé 2002b and references therein) is the most popular method for addressing the radiative transfer problem. Analytical solutions only exist for highly symmetric systems. For instance, the emerging spectrum from a plane-parallel and static homogeneous slab is characterized by two sharp peaks in the Doppler wings of the line (Neufeld 1990 and references therein). The plane-parallel solution approximately holds also for self-shielded systems where the ionized layer which surrounds the neutral region is thin with respect to the characteristic radius of the cloud. In this ideal case, optically thick systems act as efficient mirrors which convert nearly 60% of the impinging ionizing flux into Ly α photons (Gould & Weinberg 1996).

Direct imaging of fluorescent sources would lead to a major advance in our understanding of galaxy formation. Determining the size distribution of LLS at $z \gtrsim 3$ would be crucial to distinguish whether they arise from photoionized clouds in galactic halos (Steidel et al. 1995; Mo & Miralda-Escudé 1996) or in minihaloes formed prior to reionization (Abel & Mo 1998). At the same time, the intensity of the cosmic UV background could be inferred from the observed brightness of the fluorescent emission.

With present-day technology, the detection of fluorescent emission from high-redshift gas condensations is challenging, but not impossible. At $z \sim 3$, the intensity of the diffuse ionizing background (e.g. Haardt & Madau 1996) corresponds to a Ly α surface brightness of the order of $10^{-20} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2}$. It is then not surpris-

ing that blind searches have only produced a number of null results (Lowenthal et al. 1990; Martínez-Gonzalez et al. 1995; Bunker, Marleau & Graham 1998). Positive fluctuations in the ionizing background can be used to increase the signal. For instance, clouds lying close to a bright quasar are exposed to a stronger UV flux (with respect to an “average” cloud) and are then expected to be brighter in fluorescent Ly α . Very recently, Francis & Bland-Hawthorn (2004) presented a deep narrow-band search for Ly α emission in a field which lies next to the quasar PKS 0424-131. Based on quasar-absorption-line statistics and on simple models for fluorescent emission (Gould & Weinberg 1996), they expected to detect more than 6 clouds but none were seen. These null results highlight the need for a more sophisticated analysis of fluorescent Ly α emission in realistic environments.

In this paper, we present accurate models of the fluorescent Ly α emission from LLSs at redshift $z \sim 3$. Our study proceeds in three steps. First, we perform a hydrodynamical simulation of structure formation to compute the cosmological distribution of the baryons at $z = 3$. A simple radiative transfer scheme is then used to propagate the ionizing radiation through the computational box and to compute the distribution of neutral hydrogen and of recombinations. Finally, a three-dimensional Monte Carlo code is used to follow the transfer of Ly α photons. As ionizing radiation, we first consider the diffuse background generated by the UV emission of galaxies and quasars (Haardt & Madau 1996). We then discuss an inhomogeneous case where the ionizing flux from a quasar (which lies in the foreground of the gas clouds) is superimposed to the uniform background. Our detailed numerical analysis shows that simplified models (e.g. Gould & Weinberg 1996) tend to overpredict the Ly α flux emitted from optically thick regions.

The structure of the paper is as follows. We describe our numerical techniques in §2 and present our results in §3 where we also discuss the implications of our analysis for present and future observations. Finally, we discuss the limitations of our approach in §4 and we conclude in §5.

2. METHOD

2.1. Cosmological simulation

The formation and evolution of the large-scale structure in a “concordance” Λ CDM cosmological model is followed by means of an Eulerian, grid based Total-Variation-Diminishing hydro+N-body code (Ryu et al. 1993). We assume that the mass density parameter $\Omega_0 = 0.3$ (with a baryonic contribution $\Omega_b = 0.04$), the vacuum-energy density parameter $\Omega_\Lambda = 1 - \Omega_m = 0.7$ and the present-day value of the Hubble constant constant $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ with $h = 0.67$. The simulation is started at redshift $z = 60$ and follows the evolution of Gaussian density fluctuations characterized by a primordial spectral index $n = 1$ and “cluster-normalization” $\sigma_8 = 0.9$ (with σ_8 the rms linear density fluctuation within a sphere with a comoving radius of $8 h^{-1} \text{ Mpc}$). This is consistent with the most recent joint analyses of temperature anisotropies in the cosmic microwave background and galaxy clustering (e.g. Tegmark et al. 2004 and references therein). We use a comoving computational box size of $10 h^{-1} \text{ Mpc}$ where

the dark matter distribution is traced by 256^3 particles and the gas component is evolved on a comoving grid with 512^3 zones. The nominal spatial resolution for the gas (the mesh size) is $\sim 20 h^{-1} \text{ kpc}$ (comoving) with the mean baryonic mass in a cell being $\sim 10^5 h^{-1} M_\odot$. On the other hand, each dark matter particle has a mass of $5 \times 10^6 h^{-1} M_\odot$. All the results presented in this work are derived from the $z = 3$ output of a simulation which does not include radiative cooling of the gas. The limitations of this assumption are briefly discussed in §4. We defer a detailed analysis of the radiative case to future work.

2.2. Radiative transfer of UV radiation

In order to compute the distribution of neutral hydrogen within a snapshot of the computational box, we need to simultaneously solve the radiative transfer problem for UV radiation and the rate equations describing the balance between the ionization and recombination rates.

For simplicity, we assume that hydrogen is in ionization equilibrium and use the “on the spot” approximation (Baker 1962):

$$(1 - x) n_H \int_{\nu_0}^{\nu_{\text{up}}} d\nu \frac{\sigma_\nu}{h_P \nu} \int_{4\pi} d\Omega J_\nu(\Omega) = x n_H n_e \alpha_B(T) \quad (1)$$

where h_P , n_e , x , n_H , σ_ν , T and α_B respectively denote the Planck constant, the electron number density and the hydrogen ionized fraction, volume number density, ionization cross section, temperature and case B recombination coefficient (for which we use the fit by Hui & Gnedin 1997). The intensity of ionizing radiation per unit frequency and solid angle is given by J_ν (in $\text{erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ Hz}^{-1}$). The frequency integral in equation (1) extends from the hydrogen ionization threshold, $h_P \nu_0 = 13.6 \text{ eV}$, to a maximum frequency ν_{up} (which is, formally, infinite). A good approximation for our purposes is to assume $\nu_{\text{up}} = 4 \nu_0$, (i.e. set the intensity of radiation to zero at frequencies above the ionization threshold for HeII). The motivation is twofold. First, nearly all the photons with $\nu > 4 \nu_0$ (which anyway contribute only a few per cent of the energy available for H ionization in the UV background) will be absorbed by He atoms (Haardt & Madau 1996). Second, HeII recombines faster than HI and the intensity of radiation at the HeII Lyman limit is typically lower than at ν_0 . Therefore, HeII is more easily shielded from the ionizing background with respect to HI (Miralda-Escudé & Ostriker 1990). This implies that HeII-ionizing photons are absorbed in the outer regions of the gas concentrations where H is nearly fully ionized. In order to describe the hydrogen shielding layers we thus neglect HeIII and assume that the neutral fraction of He coincides with $1 - x$ (Zheng & Miralda-Escudé 2002a). For a helium abundance of $Y = 0.24$, this corresponds to assuming $n_e = \beta x n_H$ with $\beta \simeq 1.08$ (see also §2.3). Other than this, the presence of He atoms is neglected in equation (1). Given that HeII recombinations produce HI-ionizing photons and the relatively small number density of helium atoms and ions, this approximation should be reasonably accurate. Note that also recombination radiation from HeIII can ionize HI. However, considered the different spatial distribution of HeIII and HI discussed above and the characteristic HeIII-recombination time scales, we neglect the small local corrections to the HI-ionizing background deriving from this effect.

In each cell of the simulation, the diffuse ionizing background is approximately described by following the radiative transfer along 6 “light-rays” which propagate parallel (and antiparallel) to the main axes of the computational box. With this numerical trick we can treat anisotropic backgrounds (created, for instance, by shadowing effects) with a minimal request of CPU time (see Appendix A for a test of this approximation). Let us denote by $\tau_i(\nu)$ the optical depth of a given cell along the i -th ray. This quantity is computed by integrating the product $(1-x)n_{\text{H}}\sigma_{\nu}$ from a given starting location (a light source) in the box (see below) up to the first point of the cell crossed by the ray. The closest face of the cell is then exposed to a radiation field with intensity $J_{\nu,i}^{\text{in}} e^{-\tau_i(\nu)}$, where $J_{\nu,i}^{\text{in}}$ denotes the input ionizing radiation before it is filtered by the gas distribution in the box. Let us also indicate with $\Delta\tau(\nu) = (1-x)n_{\text{H}}\sigma_{\nu}L$ (with L the cell size in physical units) the optical-depth variation within the cell measured along one of its principal axes. In order to implement a photon conserving scheme, we replace the left-hand side in equation (1) with the quantity

$$\frac{4\pi}{6} \sum_{i=1}^6 \int_{\nu_0}^{\nu_{\text{up}}} \frac{J_{\nu,i}^{\text{in}}}{h\nu} e^{-\tau_i(\nu)} \frac{1 - e^{-\Delta\tau(\nu)}}{L} \quad (2)$$

where the sum is taken over the six rays (labeled by the index i). This corresponds to the number of ionizing photons (per unit volume and time) which are deposited in a given cell by the six rays. To describe the diffuse UV background, we assume that $J_{\nu,i}^{\text{in}} = J_{\nu}^{\text{HM}}$ with J_{ν}^{HM} the intensity of radiation derived at $z = 3$ by Haardt & Madau (in preparation, hereafter HM) considering the emission from observed quasars and galaxies after it is filtered through the Ly α forest¹. We assume that underdense cells are exposed to the full, isotropic background. On the other hand, overdense cells see an anisotropic radiation field which is computed by using equation (2) to propagate the input background starting from the surface where $\rho = \bar{\rho}$. The intensity of radiation (and thus x) in each overdense cell depends on the ionized fraction of the surrounding region. To solve the non-local equations, we start our calculations by assuming that the whole simulation box is optically thin (i.e. it is exposed to the input radiation field) and we iterate the radiative transfer and ionization-equilibrium calculations until convergence (within 1%) is reached in each overdense cell.

We use a similar approach to discuss the anisotropic radiation field generated by a quasar lying in the foreground of the simulation box along the observer’s line of sight. For simplicity, we assume that the quasar lies distant enough from the simulated region that its emission can be modeled as a train of plane waves impinging onto a face of the simulation box. We also assume that the quasar input spectrum is identical to that of the cosmic background. Given that J_{ν}^{HM} is well described by a power-law of index -1.25 between ν_0 and

$3\nu_0$, this is a sufficiently good approximation for our purposes (see also the extensive discussion in § 4.4). We then write the quasar ionizing flux ($\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$) as $F_{\nu} = \pi b J_{\nu}^{\text{HM}} \delta_{1i}$ with δ_{ij} the Kronecker symbol and b a dimensionless constant. This is equivalent to using $J_{\nu,i}^{\text{in}} = 1.5 b J_{\nu}^{\text{HM}} \delta_{1i}$ in equation (2). In this case, we compute the optical depth starting from the face of the simulation box which is first reached by quasar light (i.e. along the direction $i = 1$).

A self-consistent calculation of the gas temperature requires a joint treatment of radiative transfer and hydrodynamics which is still beyond present-day computing capabilities. Assuming that the photoionized gas is in thermal equilibrium, we find that $T \simeq 1 - 3 \times 10^4$ K for the typical densities in the shielding layers ($100 \lesssim \rho/\bar{\rho} \lesssim 300$). However, shock heating can easily drive the gas temperature to 10^{5-7} K. This is particularly important for the low-density regions ($\rho \lesssim 100 \bar{\rho}$) where cooling processes are inefficient and the shocked material remains hot (Theuns et al. 1998). In our analysis, we assume that $T = 2 \times 10^4$ K everywhere. This is an excellent approximation for highly overdense regions ($\rho \gtrsim 100 \bar{\rho}$) where the cooling time is shorter than the Hubble time and the gas temperature rapidly approaches the equilibrium solution (Theuns et al. 1998). Anyway, since the recombination coefficient α_{B} has only a weak dependence on T , fixing the temperature to 2×10^4 K in the whole simulation box does not seriously affect our results.

Note that, at $T = 2 \times 10^4$ K, the hydrogen recombination timescale is $t_{\text{rec}} = 2.26 (\bar{\rho}/\rho) \times 10^{10}$ yr. Ionization equilibrium will approximately hold only where t_{rec} is shorter than the characteristic quasar lifetime ($\sim 10^8$ yr, Porciani, Magliocchetti & Norberg 2004), i.e. for $\rho \gtrsim 200 \bar{\rho}$. At lower densities, our assumption of ionization equilibrium will then overestimate the hydrogen ionized fraction. This is not a problem for our study since, in the vicinity of a quasar, the ionizing flux is strong enough to nearly completely ionize the low-density intergalactic medium. It is worth noticing, however, that regions with $\rho < 200 \bar{\rho}$ will emit their recombination radiation after the quasar has switched off and will not be detectable in a survey centred onto a bright quasar.

2.3. The clumping factor

Hydro-simulations have a finite spatial resolution and cannot describe the gas distribution on arbitrarily small scales. In other words, they provide a coarse grained representation of the density field. However, the hydrogen recombination rate scales proportionally to the square of the local (i.e. fine grained) number density and is sensitive to small-scale inhomogeneities (clumpiness) within a simulation cell. In order to keep track of this discrepancy, we re-write the mean recombination rate within a cell as

$$x^2 \beta C \langle n_{\text{H}} \rangle^2 \alpha_{\text{B}}(T) \quad (3)$$

where the average is taken over a simulation cell and

$$C = \frac{\langle n_{\text{H}}^2 \rangle}{\langle n_{\text{H}} \rangle^2} \quad (4)$$

denotes the clumping factor of the gas (we assume that different atomic species and ions have the same spatial distribution). In principal, the latter quantity can be

¹ This is obtained using the most recent results regarding the quasar luminosity function and cosmic evolution within a concordance cosmological model. It assumes that the galaxy escape fraction of ionizing radiation is $f_{\text{esc}} = 0.1$ and that the energy spectral index for quasar radiation is $\alpha = 1.8$. The resulting hydrogen ionization rate is a factor 1.16 smaller than in the models by Haardt & Madau (1996) used by Gould & Weinberg (1996). The spectrum is available at <http://pitto.mib.infn.it/~haardt/refmodel.html>.

estimated by comparing simulations with different resolutions and consistent initial conditions. We assume that $\mathcal{C}$ is constant everywhere and we fix its value by imposing that the number density (per unit redshift) of LLSs in our simulation matches the observational data (Péroux et al. 2003).² This normalization procedure, which requires $\beta\mathcal{C} \simeq 6$, partially overcomes the limitations of our simulation (limited resolution and any missing physics). Note that the observational data constrain the product $\beta\mathcal{C}$ so that there is no need to specify a priori the He ionization state as discussed in §2.2.

2.4. Ly α emission

Using equation (3), we compute the hydrogen recombination rate in each cell of the simulation. In order to convert this quantity into an emission rate for Ly α photons, we need to evaluate how many recombinations ultimately lead to a $2P \rightarrow 1S$ transition. For $T = 2 \times 10^4$ K, nearly 44% of the atoms directly recombine to the ground level while 35% of the remaining cases ultimately produce excited atoms in the $2S$ state which decays to $1S$ via two-photon emission (both fractions are weakly dependent on the gas temperature, see e.g. Osterbrock 1989). Therefore, if the gas is optically thin to UV photons, only a fraction $\epsilon_{\text{thin}} = \alpha_{2P}^{\text{eff}}/\alpha_A \sim 0.36$ (where α_{2P}^{eff} and α_A denote the effective recombination coefficient to the $2P$ level and the case A total recombination coefficient, respectively) of the recombinations yield a Ly α photon. However, in the optically thick case, continuum photons produced by recombinations to the ground level can be captured by neutral atoms and produce additional Ly α radiation. The asymptotic yield in the extremely thick case (case B approximation, where no continuum photon can leave the cloud) is $\epsilon_{\text{thick}} = \alpha_{2P}^{\text{eff}}/\alpha_B \sim 0.65$. We use this value to compute the emission rate of fluorescent Ly α photons in the simulation box.

2.5. Resolving the optical depth

When we apply the method described above to our simulation, we find that the shielding layers (where the transition between optically thin and optically thick regions occurs) are poorly resolved (see Fig. 1). Typically, they consist of very few cells which each correspond to an HI optical depth variation (at the Lyman limit) of $\Delta\tau_{\text{cell}} \equiv \Delta\tau(\nu_0) \gtrsim 1$. However, for a proper treatment of the radiative transfer problem, more stringent requirements on the grid spacing must be met. In particular, the Ly α -emitting regions must be resolved with $\Delta\tau_{\text{cell}} \lesssim 1$. If not, both the spatial distribution of recombinations and the escape probabilities of Ly α photons along different directions (see §2.6) are spuriously altered.

To solve this problem, we adaptively refine the Ly α emitting regions by interpolating the original density and velocity fields of the input simulation. We use the solution of the radiative-transfer problem for the original (unrefined) grid to select the regions to interpolate and the factor of refinement. Given the memory limitations of the available machines, we use a 100^3 cells sub-box (which is particularly rich of structures) of the original




FIG. 1.— The fraction of all hydrogen recombinations happening in the simulation box which take place in cells with a single-cell HI optical depth (at the Lyman Limit) $\Delta\tau_{\text{cell}} > \Delta\tau_{\text{ion}}$ is plotted as a function of $\Delta\tau_{\text{ion}}$. The function F_{rec} equals unity when $\Delta\tau_{\text{ion}}$ is equal to (or smaller than) the minimum optical depth of a single cell in the simulation box. Dashed and solid lines respectively refer to the original and the adaptively refined simulation boxes. Note that a proper treatment of the radiative transfer problem requires that the Ly α photons are generated within cell with $\Delta\tau_{\text{cell}} \lesssim 1$ (see text).

simulation and we interpolate every cell with a significant recombination rate ($> 0.1\%$ of the maximum) and $\Delta\tau_{\text{cell}} > 1$. The level of refinement is scaled proportionally to $\Delta\tau_{\text{cell}}$ (up to a factor of 32 in each dimension) in order to have a subgrid of cells with $\Delta\tau_{\text{cell}} \lesssim 1$. Eventually, we re-compute the radiative transfer for the adaptively refined grid. Figure 1 shows that the fraction of recombinations originated in cells with $\Delta\tau_{\text{cell}} > 1$ decreases from 30% to 7% as a result of this refinement. Moreover, in the finer grid, only a negligibly small number of recombinations takes place in extremely thick cells ($\Delta\tau_{\text{cell}} > 10$) compared with 12% of the original grid.

As discussed in §2.3, we account for unresolved substructure in our simulation box by using a non-vanishing clumping factor in the equation of ionization equilibrium. Density variations within a parent cell of the original simulation due to the refinement procedure described above could, in principal, significantly contribute to the clumping factor. If this is the case, we should then adopt a value $\mathcal{C} < 6$ for the refined simulation to reproduce the observed abundance of LLSs. We find that the clumping associated with the refinement is severe in the densest zones of the simulation (which typically lie in the self-shielded regions and do not contribute to the Ly α flux) but amounts to only a few per cent in the most rapidly recombining cells. For these, we can then safely adopt $\mathcal{C} = 6$ also for the refined box.

Increasing the spatial resolution of the simulation complicates the radiative transfer of ionizing radiation generated by recombinations. Equation (1) assumes that every ionizing photon generated by a HII recombination is absorbed in the same cell in which is generated. However, this is no longer a good approximation for the adaptively refined cells which are optically thin to UV radiation. In

² Note that the spectral resolution of the observational data roughly corresponds to our box size. Therefore we can safely compute the hydrogen column density by integrating n_{HI} along the entire box.

this case, ionizing photons generated by recombinations can be absorbed in a different cell with respect to where they are created. This process is too complicated to follow without an accurate radiative transfer scheme and we use equation (1) also for the refined cells. How does this affect our results for the distribution of HI? First, the propagation of recombination radiation can slightly extend (of a few cells) the thickness of the shielding layer of a gas cloud with respect to our results. The effect is probably more pronounced in the outer shells where the gas density is lower. This should only redistribute the birth point of a small fraction of line photons. On the other hand, in the central part of the shielding layer (which contributes most recombinations) we expect that the flows of incoming and outgoing recombination radiation should nearly balance given that the hydrogen density shows little variations. In summary, our approximated treatment of recombination radiation should only slightly modify the spectral energy distribution of the emerging Ly- α line

2.6. Ly α radiative transfer

We now combine the results of the previous sections (namely, a set of arrays containing the Ly α emission rate, the HI density and the gas velocity field as a function of spatial position) to compute the spectra and the projected image on the plane of sky of the fluorescent sources. The radiative transfer of resonant Ly α photons is modeled using a three-dimensional Monte Carlo scheme analogous to that employed by Zheng & Miralda-Escudé (2002b, see also Ahn, Lee & Lee 2001). The method follows a large number of photon trajectories as they are scattered within the HI density and velocity distribution of the hydro-simulation.

2.6.1. Emission of Ly α photons

We assume that Ly α photons are isotropically emitted with frequency ν_0 in the frame of the recombining atoms (the natural linewidth is negligibly small for our purposes). In the cosmic frame (e.g. for an observer lying at the center of the simulation box and which participates to the free expansion of the universe), the frequencies of the resonant photons appear Doppler shifted by the projected velocities of the atoms along the photon trajectories. The velocity of a hydrogen atom with respect to the cosmic frame is given by the superposition of the Hubble flow with the bulk motion of the gas (i.e. the peculiar velocity of the fluid in the corresponding cell of the simulation) and a random thermal velocity:

$$\mathbf{v} = H(z)\mathbf{r} + \mathbf{v}_{\text{gas}} + \mathbf{v}_{\text{th}} \quad (5)$$

with $\mathbf{r}$ the atom position with respect to the center of the simulation box. The component of $\mathbf{v}_{\text{th}}$ along the direction of the emitted photon is generated by extracting a Gaussian deviate out of a distribution with zero mean and dispersion $\sigma_{\text{th}} = (k_{\text{B}} T/m_{\text{H}})^{1/2} = 12.8 (T/2 \times 10^4 \text{ K})^{1/2} \text{ km s}^{-1}$ (with k_{B} the Boltzmann constant and m_{H} the atomic mass).

2.6.2. Absorption

The photon frequency can be conveniently expressed in terms of the variable

$$x = \frac{\nu - \nu_0}{\Delta} \quad (6)$$

which measures the frequency shift from the Ly α line center in units of the Doppler width, $\Delta = \sqrt{2} \nu_0 \sigma_{\text{th}}/c$, where c denotes the speed of light. The mean scattering cross section of Ly α photons in the fluid frame is

$$\sigma_{\text{Ly}\alpha}(x) = \sqrt{\pi} f_{\text{Ly}\alpha} \frac{c r_e}{\Delta} H(a, x) \quad (7)$$

where $f_{\text{Ly}\alpha}=0.416$ is the Ly α oscillator strength, $r_e = 2.82 \times 10^{-15} \text{ m}$ is the classical electron radius and

$$H(a, x) = \frac{a}{\pi} \int_{-\infty}^{+\infty} \frac{e^{-y^2}}{(x-y)^2 + a^2} dy \quad (8)$$

is the Hjerting-Voigt function. For the relatively low-densities we are interested in, atomic collisions are not important and the damping coefficient a can be expressed in terms of the spontaneous decay rate Γ as $a = \Gamma/(4\pi\Delta) = 3.3 \times 10^{-4} (T/2 \times 10^4 \text{ K})^{-1/2}$.

We use equation (7) to determine the distance covered by each photon before it is scattered by an atom. We first extract a random deviate, R , from an exponential distribution function and then we integrate the product $n_{\text{HI}} \sigma_{\text{Ly}\alpha}(x)$ along the photon direction of motion until the resulting optical depth equals R . If the photon still lies within the computational volume, we select the velocity of the scatterer. Note that, in order to be able to absorb line radiation, an atom must have a velocity component along the trajectory of the incoming photon, $v_{\parallel}$, which closely matches the Doppler shift. From equation (8), it follows that, in the fluid frame, $x_{\parallel} = v_{\parallel}/(\sqrt{2}\sigma_{\text{th}})$ is characterized by the following probability distribution

$$P(x_{\parallel}) = \frac{a}{\pi H(a, x)} \frac{e^{-x_{\parallel}^2}}{(x - x_{\parallel})^2 + a^2} \quad (9)$$

We use the method presented by Zheng & Miralda-Escudé (2002b) to generate deviates which follow this statistic. The perpendicular component of the thermal velocity in the scattering plane, $x_{\perp}$, is then extracted from a Gaussian distribution with a temperature-dependent dispersion as described above.

2.6.3. Re-emission

A new direction for the photon is then randomly selected according to a phase function, $P(\cos \theta)$ (with θ the scattering angle), determined by atomic physics. Resonant scattering has an isotropic angular distribution, $P = 1$, while wing scattering is characterized by the Rayleigh phase function, $P = 3(1 + \cos^2 \theta)/4$ (Stenflo 1980). We find that the two angular distributions give consistent outputs. All the results presented in this work are obtained assuming isotropic re-emission.

To determine the new photon frequency, we assume that the scattering process is coherent in the reference frame of the scatterer (partially coherent scattering). This is appropriate when the excited atom undergoes no collisions before re-emission and the radiative damping coefficient is small (Avery & House 1968). Both conditions apply to Ly α radiation emitted by gas in the typical conditions of the shielding regions in the intergalactic medium. Once the scattering angle and the photon velocity of the scatterer are specified, it is straightforward to compute the frequency shift of the re-emitted photon in the fluid frame:

$$x = (x_{\text{in}} - x_{\parallel}) + x_{\parallel} \cos \psi + x_{\perp} \sin \psi \quad (10)$$




FIG. 2.— $\text{Ly}\alpha$ spectrum emitted by a uniform slab with a mid-plane source with optical depth τ_0 . The results of our Monte Carlo code (solid histograms) are compared with an analytical approximation (Neufeld 1990) which becomes exact in the limit $\tau_0 \rightarrow \infty$ (dotted lines). A temperature of $T = 10\text{ K}$ is assumed.

where x_{in} is the frequency shift of the incoming photon and ψ is the angle between the direction of the incident photon and the direction of the scattering atom. A Lorentz transformation is finally used to compute the frequency shift in the cosmic frame.

The set of calculations described above is iterated until the photon escapes the computational box.

2.6.4. $\text{Ly}\alpha$ spectra

To produce spectra (and broad-band images) of the fluorescent emitters, we compute the surface-brightness of the computational box along the observer's line of sight (hereafter, the x -axis). At each scattering, the probability that a photon will be re-emitted along this direction is

$$\frac{1}{4\pi} P(\cos \theta_x) e^{-\tau_x} \quad (11)$$

where θ_x is the angle between the incoming photon and the x -axis and τ_x denotes the $\text{Ly}\alpha$ optical depth of the scattering site along the observer's line of sight.³ For each photon and for each scattering, we sum this quantity to a counter in correspondence of the projected position of the scattering site and of the photon frequency. We thus obtain a three-dimensional array containing the surface brightness of fluorescent $\text{Ly}\alpha$ photons as a function of 2 spatial coordinates plus frequency. Note that a simulated photon tends to remain for many scatterings in a rather small region before it eventually escapes. This means that photons contribute only to a few pixels surrounding their emission site.

Following Zheng & Miralda-Escudé (2002b), we test our implementation of the Monte Carlo scheme against the analytical approximation by Neufeld (1990) for the optically thick, plane-parallel case. Figure 2 shows that our code accurately reproduces the analytical solution

³ This optical depth includes the effects of neutral hydrogen lying in the foreground of the computational box.

which becomes exact in the limit of extremely large optical depths.

3. RESULTS

In order to have an acceptable compromise between spectral resolution and CPU time, we only apply the Monte Carlo radiative transfer to the adaptively refined grid corresponding to a 100^3 region of the original simulation box. To achieve a good signal-to-noise ratio, we generate 10^6 photon trajectories for every simulation. We thus obtain high resolution spectra for each pixel of the resulting image that can be combined to simulate slit, line-emission integral field or broad-band observations.

3.1. Diffuse background and static gas

We first discuss the ideal case of a static gas distribution illuminated with a uniform and isotropic background of ionizing radiation. This is obtained by artificially setting to zero the velocity field of the gas within our refined box.

In the left panels of Figures 3 and 4, we respectively show the HI column density distribution and the broad-band images ($\sim 90\text{ \AA}$ in the observed frame, centered at $\lambda = (1+z) \cdot 1216\text{ \AA} = 4864\text{ \AA}$) of the selected region illuminated with the diffuse UV background. The color code in Figure 4 gives the fluorescent $\text{Ly}\alpha$ emission rate (photons per unit time, surface and solid angle) in units of the impinging rate of ionizing photons times ϵ (i.e. the fraction of the recombinations yielding a $\text{Ly}\alpha$ photon):

$$R_{\text{HM}} = \epsilon_{\text{thick}} \int_{\nu_0}^{4\nu_0} \frac{J_{\nu}^{\text{HM}}}{h\nu} d\nu = 2.44 \times 10^4 \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \quad (12)$$

with $\epsilon_{\text{thick}} \simeq 0.65$.⁴ For an observer at redshift $z = 0$, this corresponds to a $\text{Ly}\alpha$ surface brightness of

$$\text{SB}_{\text{HM}} = 3.67 \times 10^{-20} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2}. \quad (13)$$

The brightest fluorescent sources correspond to compact gas clouds with a meatball topology. This is because the diffuse UV background is bright enough to fully ionize gas concentrations with $\rho \lesssim 100 \bar{\rho}$. In general, the shielding regions either lie within virialized structures or correspond to dense gas shells which are accreting onto collapsed objects. As we will see below, the velocity field of the infalling gas produces specific signatures in the $\text{Ly}\alpha$ spectra.

The compact fluorescent sources lie along the filaments and sheets which characterize the distribution of neutral hydrogen on cosmological scales. For ease of reading, we label the three largest structures (which each have a diameter of ~ 0.4 comoving Mpc) with the letters A, B and C (see Fig. 3). Cloud C is composed of two sub-units and is a part of an elongated structure which extends towards cloud B. Similarly, a filamentary plume of gas bridges clouds A and B.

⁴ There is some observational evidence that the UV background at $z = 3$ is dominated by quasar emission with a negligible contribution from star-forming galaxies (e.g. Scott et al. 2000). In this case, the models by Haardt & Madau (in preparation) give $R_{\text{HM}} = 1.88 \times 10^4 \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. The spectral shape of the UV background between ν_0 and $4\nu_0$ is nearly identical to the general case discussed in the main text. Therefore, our predictions for the surface brightness of fluorescent sources can be simply scaled down by 30% if future observations will prove that galaxies do not significantly contribute to the ionizing background at $z = 3$.




FIG. 3.— Column-density distribution of neutral hydrogen at $z = 3$. In the left panel, the gas is exposed to a diffuse UV background generated by the population of galaxies and quasars. In the right panel, the ionizing flux from a foreground quasar, located a short distance in front of the region and corresponding to a boost factor $b = 6$ (see equation (14)) is superimposed to the diffuse background.




FIG. 4.— Broad band images ($\sim 90 \text{ \AA}$, centered on $4864 \text{ \AA}$) of fluorescent Ly α emission at $z = 3$ for static gas clouds (left) and accounting for the gas velocity field (right). Both images correspond to the column density distribution presented in the left panel of Figure 3. The white boxes indicate the location of the slit spectrographs used to obtain the energy distributions presented in Figure 7.

Simple reasoning based on the plane-parallel model for line transfer suggests that, in the absence of photon sinks (e.g. dust), self-shielded (isotropically-illuminated) objects should shine with a surface brightness of SB_{HM} (Hogan & Weymann 1987; Gould & Weinberg 1996). In our static simulation (Fig. 4, left panel), the SB of self-shielded objects closely matches the predictions of this simple plane-parallel model. The SB distribution in the simulation (dotted histogram in Fig. 5) shows a narrow peak at this expected value. In general, the SB scales proportionally to $N_{\text{HI}}^{1/2}$ in the optically thin regions and asymptotically approaches its maximum value for self-shielded objects (see the top-left panel in Fig. 6). The brightest lines of sight in fluorescent Ly α corre-

spond to optically thick systems with column densities $N_{\text{HI}} \gtrsim 10^{18} \text{ cm}^{-2}$ which are thus associated with LLSs and DLAs. All the photons of the ionizing background are converted into Ly α radiation within the shielding layers of these optically thick systems. In the absence of other sources of ionizing radiation, it is impossible to produce a stronger Ly α flux. This explains why the brightest objects in the left panel of Figure 4 have a uniform SB and sharp boundaries which correspond to the regions with $N_{\text{HI}} \simeq 10^{18} \text{ cm}^{-2}$ in the left panel of Figure 3.

3.2. Diffuse background and realistic gas velocities

We are now ready to discuss the more realistic case where we include the gas velocity field of the hydro-




FIG. 5.— Surface brightness distribution of fluorescent sources ionized by a diffuse UV background (solid), by the additional contribution of a quasar with “boost” factor $b = 2$ (dashed) and $b = 6$ (dot dashed). The dotted line is analogous to the solid one but is obtained by artificially setting to zero the gas velocity field.

simulation. The corresponding Ly α emission rate is shown in the right panel of Figure 4. The overall pattern is similar to the static case, but a number of striking differences are noticeable. Namely: *i*) the SB of self-shielded objects is no longer uniform (e.g. the right-hand side of Cloud A is nearly a factor of 2 fainter than the left-hand side); *ii*) the boundaries of the emitting regions are less sharp and self-shielded objects are surrounded by large, low-SB halos; *iii*) self-shielded objects can be significantly fainter (or, very rarely, brighter) than in the static case.

The top-right panel in Figure 6 shows that the gas velocity field introduces additional scatter into the SB - N_{HI} relation with respect to the static case. The brightest lines of sight still correspond to $N_{\text{HI}} \gtrsim 10^{18} \text{ cm}^{-2}$ but now two regions with the same column density can be associated with brightnesses which differ up to a factor of 5. In consequence, the SB distribution of optically thick regions is broader and it is slightly shifted to fainter fluxes with respect to the static case (see the peak of the solid histogram in Fig. 5). We find that the median SB of the self-shielded objects amounts to nearly 75% of the value predicted by Gould & Weinberg (1996). At the same time, a larger fraction of the sky has $\text{SB} < \text{SB}_{\text{HM}}$ compared to the static case. (the power-law part of the solid histogram in Fig. 5). As we will show below, this excess is caused by foreground scattering of the Ly α photons and is related to the presence of extended Ly α halos around self-shielded objects.

A better understanding of the “velocity-field effect” can be achieved by comparing the spectra of the fluorescent emission in the static and in the general case. In the left and central panels of Figure 7, we show the corresponding spectral energy distributions of the Ly α photons. These have been obtained positioning four slit spectrographs (width $\simeq 0.9$ arcsec and variable length) on top of the three brightest sources as shown in the right panel of Figure 4. In a static gas distribution, spectra




FIG. 6.— The Ly α surface brightness of each pixel of the simulated images is plotted against the corresponding column density of neutral hydrogen. In the top panels, the intergalactic medium is ionized by a diffuse background. In particular, the top-left frame refers to a static gas distribution. In the bottom panels, a quasar with boost factor $b = 2$ (bottom-left) and $b = 6$ (bottom-right) is superimposed to the diffuse background. Dotted lines mark the expected SB for a plane-parallel slab while dashed lines indicate the minimum column density for LLSs (short-dashed) and DLAs (long-dashed).

have a characteristic double humped shape and are symmetric with respect to the line center. On the other hand, in the general case the energy distribution is no longer symmetric. In fact, particular configurations of the velocity and density fields are able to strongly suppress one of the wings of the Ly α line and significantly lower the observed SB of the self-shielded objects. In the particular case of Cloud A, a low-density concentration of neutral hydrogen is infalling onto the Ly α emitting region. The relative velocity (along the line of sight) corresponds to $\sim 4\sigma_{\text{th}}$ and thus to a very high optical depth. Therefore, most of the photons that, in the static case, leave the shielding layers along the line of sight in the red Doppler wing will scatter within the infalling cloud and escape in other directions lowering the observed surface brightness. These photons will then form the extended Ly α halos which surround the brightest objects in Figure 4. The phase-space distribution of neutral gas in the vicinity of the emitting regions thus plays a fundamental role in reshaping the Ly α spectral energy distribution. In broad terms, infalling material diminishes the red wing of the spectrum, while gas which is receding from the emitting region (which could also mean that the shielding layer is infalling onto a central object more rapidly than the surrounding gas) damps out the blue peak of the spectrum. We find that the velocity dispersion of the gas within the regions crossed by the Ly α photons broadens the red and blue peaks of the spectrum by up to 10 Å. On the other hand, when both peaks are detectable, their separation is nearly independent from the detailed properties of the emitting regions and is set by the thermal velocity dispersion of the original cloud. In fact, in analogy with the plane-parallel case, the escape probability of a Ly α photon peaks at a frequency which only depends on




FIG. 7.— Two-dimensional spectra obtained “observing” our simulations with the slits shown in the right panel of Figure 4. Different columns refer to different simulations. In particular, from left to right: diffuse background and static gas distribution (§3.1), diffuse background and realistic gas velocities (§3.2), quasar with $b = 6$ plus diffuse background (§3.3). The labels in the top left corner indicate a particular slit spectrograph as in Figure 4.

the optical depth of its emission site and on the temperature of the medium. For a typical self-shielded cloud ($\tau_{LL} \gtrsim 1$), the spectrum peaks at $\sim \pm 4x$ which, for the assumed temperature, corresponds to a separation of ~ 8 Å.

The two-dimensional spectra shown in the central panel of Figure 7 clearly show that the gas velocity field within and in the vicinity of the shielding layers has a complicated structure which does not show the characteristic pattern of ordered rotation or symmetric infall considered by Zheng & Miralda-Escudé (2002b).

3.3. Quasar plus diffuse background

We now discuss a case of anisotropic illumination, obtained by superimposing the ionizing flux from a quasar to the diffuse UV background. The quasar is imagined to lie a short distance in front of the computational box as seen by us, and thus enhances the UV illumination experienced by faces of gas clouds exposed to it. Note that the “boost” factor b (defined in §2.2) is determined by the intrinsic luminosity of the quasar and by its actual separation from the simulated region. The definition above can be generalized to any given quasar spectrum using the emitted rate of ionizing photons. At a physical distance r from a quasar with monochromatic luminosity

$L_\nu = L_{LL}(\nu/\nu_{LL})^{-\alpha}$, we find

$$b = 15.2 \frac{L_{LL}}{10^{30} \text{ erg s}^{-1} \text{ Hz}^{-1}} \frac{0.7}{\alpha} \left(\frac{r}{1 \text{ Mpc}} \right)^{-2}. \quad (14)$$

The resulting N_{HI} distribution (assuming a boost factor $b = 6$) is shown in the right panel of Figure 3. The corresponding broad-band image (obtained accounting for gas velocities) is presented in the left panel of Figure 8. As expected, the self-shielded regions (and thus the fluorescent sources) are smaller with respect to the isotropic background case due to the extra-ionizing radiation produced by the quasar. This also makes the fluorescent sources brighter (dot-dashed histogram in Fig. 5) since more recombinations will be produced to balance a stronger ionization rate. Based on the (plane-parallel) slab model, where Ly α photons are emitted following a cosine law (Gould & Weinberg 1996), one would have naively expected an increase in the Ly α surface brightness towards the observer by a factor $1 + b = 7$ with respect to the diffuse background case.⁵ However, Figures 5 and 6 clearly indicate that the slab model overestimates

⁵ This holds for normal incidence. In general, the surface brightness of a slab which forms an angle θ with the incident quasar flux corresponds to a factor $1 + b \cos \theta$.




FIG. 8.— Broad band images ($\sim 90 \text{ \AA}$, centered on $4864 \text{ \AA}$) of fluorescent $\text{Ly}\alpha$ emission from our simulation box at $z = 3$. In both cases, the intergalactic medium is ionized by a diffuse background and by a quasar with boost factor $b = 6$. The image on the left (right) is obtained by observing our simulation box from a line of sight parallel (perpendicular) to the direction of quasar illumination. The left frame corresponds to the column density distribution presented in the right panel of Figure 3.

the SB of the self-shielded objects. This is not due to shadowing effects. In fact, the attenuation of the quasar flux by diffuse gas lying in front of the fluorescent clouds is generally negligible. Comparing with a static simulation, we also find that gas motions can only explain a small part of this discrepancy. In fact, in the presence of a quasar, foreground scattering is reduced due to the lower neutral fraction present in low density gas and broad-band images tend to be more uniform than in the case of isotropic illumination. On the other hand, the slab approximation no longer applies when the size of the shielding layers is comparable with the radius of a cloud. In this case, $\text{Ly}\alpha$ photons produced at a particular point leave the cloud with a different angular distribution with respect to the plane-parallel case. For approximately spherical clouds and in the presence of uniform illumination, this effect is suppressed for symmetry reasons. However, when the ionizing flux is anisotropic, the $\text{Ly}\alpha$ SB does depend quite strongly on the geometry of the shielding layers.

To study how the SB of self-shielded objects along the quasar direction, $\text{SB}_{\parallel}$, depends on the impinging flux, we performed a series of simulations with increasing b . Our results are summarized in Figure 9, where we express $\text{SB}_{\parallel}$ in terms of an “effective boost factor” defined by

$$\text{SB}_{\parallel} = (1 + b_{\text{eff}}) \text{SB}_{\text{HM}}. \quad (15)$$

Points with errorbars mark the 25th, 50th and 75th percentiles of $1 + b_{\text{eff}}$ among the DLAs. The solid line represents the best-fitting relation

$$1 + b_{\text{eff}} = 0.74 + 0.50 b^{0.89}, \quad (16)$$

while the dashed line shows the predictions of the slab model. Note that the geometric effect becomes more and more important as b is increased.

Where do the “missing” $\text{Ly}\alpha$ photons go? In the right panel of Figure 8, we show the fluorescent emission along

a line of sight perpendicular to the direction of quasar illumination (assuming $b = 6$ as in the left panel). In this case, the plane-parallel model predicts that the self-shielded objects should emit at SB_{HM} . In our simulations, however, the shielding layer deeply penetrates in the clouds along the quasar direction and the slab model does not apply. In consequence, self-shielded objects are much brighter than a slab along this line of sight. Typically, $\text{SB}_{\perp} \simeq 0.5 \text{SB}_{\parallel}$ for $b \gg 1$ while $\text{SB}_{\perp} \simeq \text{SB}_{\parallel}$ for $b \ll 1$. In other words, $\text{Ly}\alpha$ photons generated by the quasar ionizing flux are emitted within a wide solid angle. As a consequence of this partial isotropization, self-shielded clouds are fainter than expected (based on the slab approximation) along the quasar direction and brighter in the perpendicular directions.

Finally, in the bottom panels of Figure 6, we show the SB - N_{HI} scatterplot for anisotropic illumination (when observer, quasar and the simulation box are aligned). It is worth noticing that, while the SB keeps nearly constant for LLSs, on average, it steadily increases with N_{HI} for DLAs. This phenomenon can be explained by as follows. Let us assume that self-shielded objects are nearly spherically symmetric. Then, i) the ionizing flux from the quasar depends on the incident angle with respect to the local density gradient in the clouds; ii) this cosine approaches unity for the central projected regions of self-shielded objects; iii) the column density reaches the highest values along these lines of sight.

3.4. Size distribution of $\text{Ly}\alpha$ sources

Knowing the size distribution of fluorescent $\text{Ly}\alpha$ sources is fundamental to planning an observational campaign for their detection. Regrettably, our refined box is too small (its size being $\sim 2 h^{-1}$ comoving Mpc) to provide a statistically representative sample of optically thick sources. On the other hand, performing the line transfer on the $10 h^{-1}$ Mpc box would require an excessive amount of computer time. For these reasons, we




FIG. 9.— Ly α surface brightness of optically thick clouds [expressed in terms of the effective boost factor defined in equation (15)] as a function of the impinging quasar ionizing flux [expressed in terms of the quasar boost factor, b , defined in equation (14)]. Points with errorbars denote the 25th, 50th and 75th percentiles of b_{eff} for the DLAs. The solid line represents the best-fitting relation given in equation (16). Predictions of the static, plane-parallel model are plotted with a dashed line.

decided to propagate only the ionizing radiation through the $10 h^{-1}$ Mpc box and to use the scatterplots in Figure 6 to convert the neutral-hydrogen column densities into Ly α fluxes. In fact, independently of the value of b , all lines of sight with $N_{\text{HI}} > 10^{18} \text{ cm}^{-2}$ are approximately associated with a constant Ly α surface brightness (within a factor of 2 uncertainty caused by the gas motion and cosine effects discussed above). We then adopt this threshold value to derive the size distribution of fluorescent objects. In Figure 10, we present our results for an isotropic ionizing background ($b = 0$). Solid and dashed histograms respectively refer to objects with $N_{\text{HI}} > 10^{18} \text{ cm}^{-2}$ and to DLAs. It is worth remembering that we fixed the value of the clumping factor in our simulation so as to reproduce the observed sky covering factor of LLSs. In consequence, if a significant fraction of the real systems have a characteristic size which is smaller than our numerical resolution, our simulation will overpredict the number of large systems in order to preserve the required normalization.

In Figure 11, we plot the number density of self-shielded objects as a function of b . We use three different thresholds for the source size: 3 (which corresponds to barely resolved objects), 20 and 80 arcsec². In all cases, the number of sources rapidly drops with increasing b . In fact, higher values of b characterize regions which are closer to a given quasar (see eq. (14)) and, obviously, correspond to a lower number density of self-shielded objects.

From this figure, it is also possible to determine the number density of sources which are brighter than a certain threshold value $(1 + b_{\text{min}}) \text{SB}_{\text{HM}}$. Let us consider a Ly α source which is optically thick to ionizing radiation at a given distance from a quasar. Let us also imagine that we can move the cloud towards the quasar




FIG. 10.— Differential size distribution of optically thick clouds (solid histogram) and DLAs (shaded histogram) in a simulation where the intergalactic medium is exposed to a diffuse ionizing background. The shaded histogram has been slightly shifted in the horizontal direction to improve readability.




FIG. 11.— Physical number density of fluorescent Ly α sources (with size A indicated by the labels) as a function of the impinging quasar flux [expressed in terms of the boost factor b defined in equation (14)]. Errorbars are derived assuming Poisson statistics. Triangles and circles have been slightly displaced in the horizontal direction to improve readability.

thus increasing the b factor. As long as the cloud keeps optically thick, $\text{SB}_{\parallel}$ monotonically increases. However, there exists a particular value of the boost factor, b_{ss} , at which the cloud is no longer able to self-shield. Therefore, for $b \gtrsim b_{\text{ss}}$, $\text{SB}_{\parallel}$ keeps roughly constant.⁶ Thus, the number of self-shielded objects at a given b coincides with the number of sources (which are not necessarily optically thick) with $\text{SB} \gtrsim [1 + b_{\text{eff}}(b)] \text{SB}_{\text{HM}}$. In other

⁶ The fraction of recombinations yielding a Ly α photon decreases from $\epsilon_{\text{thick}} \sim 0.65$ to $\epsilon_{\text{thin}} \sim 0.36$ when a cloud becomes optically thin. Therefore, we expect a fully ionized cloud to be a factor of ~ 2 fainter in Ly α with respect to the optically thick case.

words, the number of sources which are brighter than a given threshold can be computed with the following procedure. First, convert the threshold SB into an effective boost factor, $b_{\text{eff}}^{\text{thr}}$. Second, invert equation (16) to find the value of b_{min} such that $b_{\text{eff}}(b_{\text{min}}) = b_{\text{eff}}^{\text{thr}}$. Third, use $b = b_{\text{min}}$ in Figure 11 to determine the number density of the sources. Fourth, use equation (14) to find the volume within which it is possible to have $b > b_{\text{min}}$.

The variation of the number density of fluorescent sources as a function of b is somehow related to the proximity effect. Hydrogen clouds in the vicinity of a quasar are strongly ionized and emit fluorescent radiation. In other words, the missing absorption systems which determine the proximity effect can be detected in emission through their recombination radiation. Therefore, in analogy with studies of the proximity effect (e.g. Bajlik, Duncan & Ostriker 1988; Scott et al. 2000), the number density variation of fluorescent sources around a quasar can be used to infer the intensity of the UV background. In this case, reliable models of fluorescent emission are fundamental to convert the observed counts into a background amplitude.

3.5. Comparison with recent observational data

We can use the above to compare the predictions of our models with the recent observational results by Francis & Bland-Hawthorn (2004, hereafter FBH). These authors performed a deep narrow-band search for fluorescent Ly α emission in the vicinity of the $z = 2.168$ quasar PKS 0424-131 ($L_{\text{LL}} = 1.67 \times 10^{30} \text{ erg s}^{-1} \text{ Hz}^{-1}$, $\alpha \simeq 0.7$). At the 5σ confidence level (corresponding to a surface brightness of $4.7 \times 10^{-19} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2}$ for sources larger than 100 arcsec^2 and to $9.6 \times 10^{-18} \text{ erg cm}^{-2}$ for unresolved sources) no source was been detected. Based on the observed abundance of LLSs, FBH expected to find ~ 6 fluorescent clouds with a size of 100 arcsec^2 . This estimate, however, does not take into account the ionizing radiation from the quasar.

Assuming that our results at $z = 3$ are approximately valid at the quasar redshift,⁷ we find that the sensitivity limits of FBH correspond to $b_{\text{min}} \sim 11.2$ for sources which are larger than 100 arcsec^2 . Assuming that the ionizing background keeps roughly constant in the redshift interval $2 < z < 3$, from equation (14) we find that this corresponds to a distance from the quasar of $r_{\text{max}} \sim 1.5$ (physical) Mpc. This is the maximum distance from the quasar at which an optically thick cloud could have been detected. Based on our simulations, we expect to find, on average, $1 - 2$ objects within a sphere of radius r_{max} around the quasar. However, FBH limited their search to distances smaller than 1 Mpc thus reducing the sampled volume by a factor 3.4 with respect to the theoretical limit. In this case, the expected number of sources ranges between 0.3 and 0.6. Therefore, the probability of detecting (at least) one source with one single observational run is $0.25 < P < 0.45$ (assuming Poisson statistics). Our simulations clearly show that the results by FBH are thus perfectly consistent with our understanding of the intergalactic medium at high- z .⁸

⁷ We simply assume that the Ly α surface brightness scales as $(1+z)^{-4}$, i.e. $\text{SB}(z=2.168) = 2.54 \text{ SB}(z=3)$.

⁸ The detection limit for unresolved sources is less interesting.

There are caveats to the simple calculations described above. For instance, we have assumed that all the Ly α sources lying within a distance r_{min} from the quasar are brighter than b_{min} . This assumption tends to overestimate the number of detectable objects. In fact: i) self-shielded clouds lying in front of the quasar are much fainter and hardly detectable (their SB actually depends on the angle between the line of sight and the quasar direction); ii) fully ionized clouds in the foreground of the quasar tend to be a factor of $\epsilon_{\text{thick}}/\epsilon_{\text{thin}} \simeq 1.8$ fainter than assumed above; iii) if the age of the quasar is shorter than the hydrogen recombination timescale no fluorescent sources will be detectable (see the discussion at the end of §2.2). On the other hand, we have assumed that our simulation box is representative of the gas distribution surrounding a quasar. Since optically selected quasars at high- z tend to sit within the most massive dark-matter halos formed at that epoch (Porciani et al. 2004), it is reasonable to expect that matter clusters (and moves with larger peculiar velocities) around them. Therefore, we could have underestimated the number densities of fluorescent sources lying close to a quasar.

Despite of the approximations listed above, we believe that our results provide the most reliable estimates for the abundance of fluorescent Ly α sources at high- z carried out to date. Optimized sampling strategies are certainly required to observe these objects. Our simulations then represent a fundamental tool to plan observations around a given quasar.

4. UNCERTAINTIES

4.1. Radiative cooling

While numerical simulations are a useful tool to guide our understanding, they cannot be considered a perfect model of reality. A potential limit of our simulation is the lack of radiative cooling. While this is not a concern for the diffuse intergalactic medium, it becomes worrisome for highly overdense regions. Fluorescent Ly α sources have intermediate overdensities ($\rho/\bar{\rho} \sim 200$) and are likely to be in equilibrium with the UV ambient radiation. We thus expect them to be mildly affected by cooling processes. In any case, most of the results discussed in this paper are nearly independent of the details of the gas distribution. Both the velocity and the geometric effects will be present anyway. On the other hand, the size distribution of the sources might be more affected by the cooling processes. It is also worth stressing that there is no way of accounting for the effects of cooling and heating in a self consistent way. In fact, simulations where radiative transfer is fully coupled with hydrodynamics are still not viable with current supercomputers. Moreover, other poorly understood processes (like energy and momentum feedback) play an important role. Therefore, it is hard to judge the level of approximation that current simulations including gas cooling may reach.

4.2. Resolution and sub-structure

Is the finite resolution of our simulation affecting our results? Multiple metal lines are often associated with

It corresponds to $b_{\text{min}} \sim 388$ (assuming that equation (16) can be extrapolated to such high values of b) and $r_{\text{max}} \sim 250 \text{ kpc}$. The number of expected sources in the associated volume of $\sim 0.06 \text{ Mpc}^3$ is therefore negligibly small.

single DLAs (e.g. Prochaska & Wolfe 1997) thus suggesting the presence of a clumpy medium. Unresolved sub-structure in our simulation might reduce the velocity effect and modify the outcoming spectra. The adopted value of $\mathcal{C}$ implies that at least 1/6 of the volume is in dense clumps. If these sub-structures have a diameter which is comparable to the cell size, we only expect a minor modification of our results. In fact, the gas velocity in the simulation should closely approximate the motion of these clumps. The only effect is then a slight Doppler-shift of the entire Ly α spectrum of each cell. The opposite case, where each cell contains a large number of small substructures (Abel & Mo 1998), can be approximately discussed by considering an additional contribution to the thermal velocity dispersion. Assuming a value of $\sigma_{\text{th}} \sim 50 \text{ km s}^{-1}$ (corresponding to roughly half the virial velocity of the host halos, Haehnelt, Steinmetz & Rauch 1998), we find that the velocity effect is be strongly suppressed and the spectra are in better agreement with the slab model. Note, however, that the existence of a sea of small subclumps is disfavored by observations. In fact, this scenario would produce broad absorption features instead of the multiple metal systems associated with single DLAs (cf. Haehnelt, Steinmetz & Rauch 1998; McDonald & Miralda-Escudé 1999). In any case, the velocity dispersion within our simulated DLAs is of the order of 100 km s^{-1} , in good agreement with observational data.

4.3. Additional sources and dust

Beyond fluorescent emission, additional Ly α radiation might be produced in the inner regions of the clouds. Within gravitationally collapsed objects, the gas tends to dissipate its internal energy by emitting line photons (e.g. Haiman, Spaans & Quataert 2000; Fardal et al. 2001; Furlanetto et al. 2005). Similarly, internal star formation could act as a copious source of Ly α photons, but, whereas fluorescent emission is expected to extend over several tens of kiloparsec, the Ly α emission from star forming region should be more concentrated near the centers of galaxies (Furlanetto et al. 2005). We have focused here on the fluorescent emission generated by recombinations and these extra sources of line photons are not considered in our analysis. We will present a comprehensive model of Ly α emitters in a future paper.

At the same time we did not consider the destruction of Ly α photons by dust grains. Little is known about the dust properties within the intergalactic medium at $z \sim 3$ even though there is some evidence for the presence of dust in DLAs (Fall & Pei 1993). Nevertheless, the associated absorption of fluorescent photons is likely to be minimal due to the relatively low N_{HI} of the shielding layer (e.g. Gould & Weinberg 1996). On the other hand, absorption is expected to be more severe for Ly α photons produced close to and within the star-forming regions where dust is likely to be more abundant and the Ly α escape probability is lower.

4.4. The UV background

Our results are based on the Haardt-Madau model for the UV background. At $z = 3$, the adopted model has an amplitude at the Lyman limit of $J(\nu_0) = 4.04 \times 10^{-22} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ Hz}^{-1}$ and corresponds to an hydrogen ionization rate $\Gamma = 1.15 \times 10^{-12} \text{ s}^{-1}$. Even though on the low

side, this is consistent with recent observational studies of the proximity effect, which give $J(\nu_0) = 7.0^{+3.4}_{-4.4} \times 10^{-22} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ Hz}^{-1}$ and $\Gamma = 1.9^{+1.2}_{-1.0} \times 10^{-12} \text{ s}^{-1}$ (Scott et al. 2000). Our results should then considered conservative, because they account for a UV background intensity which is a factor of 1.75 lower than current observational estimates. Note, however, that a different normalization of the UV background cannot change our predictions for the number density of fluorescent sources. In fact, we determined the clumping factor of the gas by imposing that the mean number of LLS in the simulation matches the observational data. Therefore, a stronger ionizing radiation field would require a larger clumping factor to fit the observed counts. If both J_{ν}^{HM} and $\mathcal{C}$ are multiplied by the same multiplicative factor, A_J , the solution of the ionization equilibrium does not change. In fact, this is equivalent to multiply both sides of equation (1) by a constant factor. Therefore our results for the number density and size distribution of fluorescent sources are robust with respect to the amplitude of the UV background. Note, however, that the Ly α surface brightness of optically thick cloud would be higher by a factor A_J with respect to equation (13). Consistently, the boost factor in equation (14) would decrease by a factor A_J while equation (16) would remain unaltered.

Some (most likely minor) modification of our number counts is instead expected if the spectral energy distribution of the UV background significantly differ from the assumed one. This crucially depends on the relative contribution of galaxies and quasars to the cosmic ionizing background. Similarly, the spectral energy distribution of quasar radiation plays an important role. For simplicity, we assumed that the quasar spectrum is identical to that of the cosmic background. This corresponds to a energy index $\alpha \simeq 1.25$ i.e. to a much flatter spectrum than inferred from quasar observations ($1.5 \lesssim \alpha \lesssim 1.8$). What could be the effect of this assumption? The average mean free path of ionizing photons from a spectrum with $\alpha = 1.8$ is only a factor of 20% smaller than for $\alpha = 1.25$. Therefore, radiation from a steep quasar spectrum should penetrate a bit less within hydrogen clouds with respect than our models. This might slightly modify the emerging Ly α spectra and reduce the importance of the geometric effect. Note, however, that quasar radiation will be partially filtered by the IGM before reaching the fluorescent clouds. Thus, a spectrum with $\alpha = 1.8$ at emission will be transformed into a flatter distribution in the shielding layer of a cloud.

5. SUMMARY

We have presented a new method to produce realistic simulations of fluorescent Ly α sources at high redshift. We started by simulating the formation of baryonic large-scale structure in the Λ CDM cosmology. A simple radiative transfer scheme was then used to propagate ionizing radiation through the computational box and to derive the distribution of neutral hydrogen. Finally, the transport of Ly α photons generated by hydrogen recombinations was followed using a three-dimensional Monte Carlo code. As ionizing radiation, we first considered the smooth background generated by galaxies and quasars. Then, as a second case, we superimposed to the background the UV flux produced by a quasar lying in the vicinity of the simulation box.

Our detailed numerical treatment improves upon previous work which was either based on rather crude approximations for the transfer of resonantly scattered radiation (Hogan & Weymann 1987; Gould & Weinberg 1996) or on highly symmetric semi-analytical models for the gas distribution (Zheng & Miralda-Escudé 2002b). Our results show that simple models (e.g. Gould & Weinberg 1996) tend to overpredict the Ly α flux emitted from optically thick clouds. In fact, we identified two effects that reduce the fluorescent Ly α flux (and modify the spectral energy distribution) with respect to the widespread static and plane-parallel model.

Velocity effect – The velocity field inside and around the shielding layers of a gas cloud influences the emerging line profile. The symmetry of the double humped spectrum is lost and, in most cases, one of the two peaks is severely suppressed. On average, the SB of a cloud is reduced by 25% with respect to the static situation.

Geometric effect – For anisotropic illumination and in the presence of a strong ionizing flux, the thickness of the shielding layer is comparable to the size of the gas cloud. In this case, the angular distribution of the emerging radiation is very different than in the plane-parallel approximation. For instance, close to a quasar, a cloud emits much less than predicted by the slab model in the direction of the quasar and much more in the other di-

rections.

The importance of these effects (in particular of the angular redistribution of Ly α photons) depends on the intensity of the impinging radiation. In equation (16), we provided a fitting function for the maximum Ly α brightness of optically thick sources as a function of the incident ionizing flux. In Figure 11, we presented our predictions for the number density of fluorescent sources with different sizes.

These results are consistent with the recent null detection by Francis & Bland-Hawthorn (2004) and represent a fundamental tool for planning future observations.

We would like to thank Piero Madau for suggesting to consider the problem of anisotropic illumination in the presence of a quasar. We benefited from extensive discussions with Francesco Haardt, Martin Haehnelt, Tom Theuns and all the participants to the “Galaxy-Intergalactic Medium Interactions” program recently held at the Kavli Institute for Theoretical Physics in Santa Barbara. We thank the anonymous referee for useful suggestions which improved the presentation of our work. CP and FM acknowledge support from the Zwicky Prize Fellowship program at ETH-Zürich and SC from the Swiss National Science Foundation.

APPENDIX

RADIATIVE TRANSFER FOR A DIFFUSE BACKGROUND

The fluorescent Ly α flux from a source is proportional to the number of hydrogen ionizations taking place in the corresponding gas cloud. In order to simplify the radiative transfer problem for continuum radiation, in equation (2) we only followed the propagation of ionizing photons along the 6 principal directions of the simulation box. In this appendix, we first provide justification for the multiplicative factor $2\pi/3$ which appears in equation (2) and then test the 6-direction approximation.

The factor $2\pi/3$ in equation (2) is obtained by discretizing the left-hand side in equation (1) along 6 preferred directions. Note, however, that equation (1) applies to an infinitesimal volume element while our discretized version is used to describe finite cubic cells. It is clear that the approximation is correct for optically thin cells, but where does it break down? Let us consider a homogeneous cube of side L (corresponding to an optical depth $\Delta\tau_{\text{cell}}$) illuminated by a uniform and monochromatic background with intensity I (photons per unit time, surface and solid angle). The ionization rate within the cube generated by the radiation background impinging on one face can be written as $\alpha \pi I L^2 e^{-\Delta\tau_{\text{cell}}}$, with α a numerical coefficient of order unity. We use a Monte Carlo method to compute α as a function of $\Delta\tau_{\text{cell}}$. Our results are plotted in Figure 12. When the cell is optically thin $\alpha \simeq 2/3$, while $\alpha \rightarrow 1$ when $\Delta\tau_{\text{cell}} \rightarrow \infty$. A rather sharp transition between the two asymptotic regimes is observed for $\Delta\tau_{\text{cell}} \sim 10$. Note that, in the presence of a uniform background, the optically thin approximation adopted in the main text is accurate even for $\Delta\tau_{\text{cell}} = 1$, where $\alpha = 0.70$. Given that, in our adaptively refined cosmological simulation, $\sim 93\%$ of recombinations take place in cells with $\Delta\tau_{\text{cell}} < 1$, we used $\alpha = 2/3$ in our calculations.

All this, however, applies to an isolated cell exposed to the ionizing background. We now want to test how accurately equation (2) works in the shielding layer of an optically thick cloud. We thus consider a spherically symmetric cloud which is optically thin at his external boundary and thick at the center ($\log \Delta\tau_{\text{cell}} = 2 - 4(r/R)$, with r the radial coordinate which vanishes at the cloud center and R the characteristic size of the cloud, which roughly mimics one of the clouds in our refined simulation). We then illuminate the outer boundaries of the cloud with a uniform (over the external 2π sr) and monochromatic background. This is injected from the surface of a cube of side $2R$ centered on the cloud and divided into 151^3 mesh points. In Figure 12, we compare the outcome of a detailed radiative transfer code (Porciani & Madau, in preparation) with our approximated method. Note that in our six-directions approximation the ionizing flux penetrates slightly deeper into the cloud with respect to the exact solution. This happens because our method ignores photon trajectories which are oblique to the principal axes of the box. On the other hand, we find that our approximation produces 92% of the ionizations that take place in the cloud. Given the simplicity of our algorithm, this is a remarkable achievement. Note that, adopting the optically thick approximation ($\alpha = 1$), we would have overestimated the number of recombinations by 38%. These figures are rather stable and, for realistic cases, do not depend on the details of the optical depth distribution.

REFERENCES

- Abel, T. & Mo, H. J. 1998, ApJ, 494, L151
- Ahn, S., Lee, H., & Lee, H. M. 2001, ApJ, 554, 604




FIG. 12.— Left: The ionization rate generated by a uniform UV background (of intensity I) penetrating in a cubic cell (of size L) through one of its faces as a function of the optical depth variation within the cell. The solid line represents the ionization rate (in units of $\pi I L^2 e^{-\Delta \tau_{\text{cell}}}$) while the dashed line marks the optically thin solution $\alpha = 2/3$. Right: Radial profile of the ionization rate (arbitrary units) for the mock cloud described in Appendix A. The exact solution (solid) is compared with our six-direction approximation in the optically thin (dashed) and optically thick (dotted) cases.

- Avery, L. W., & House, L. L. 1968, *ApJ*, 152, 493
 Bajtlik, S., Duncan, R. C., & Ostriker, J. P. 1988, *ApJ*, 327, 570
 Baker, J., & Menzel, D. 1962, in *Selected Papers on Physical Processes in Ionized Plasma*, ed. D. Menzel (London: Constable & Co), 58
 Bunker, A. J., Marleau, F. R., & Graham, J. R. 1998, *AJ*, 116, 2086
 Cen, R., Miralda-Escudé, J., Ostriker, J. P., & Rauch, M. 1994, *ApJ*, 437, L9
 Fall, S. M., & Pei, Y. C. 1993, *ApJ*, 402, 479
 Francis, P. J. & Bland-Hawthorn, J. 2004, *MNRAS*, 353, 301
 Furlanetto, S. R., Schaye, J., Springel, V., & Hernquist, L. 2005, *ApJ* in press, astro-ph/0409736
 Gould, A. and Weinberg, D. H., 1996 *ApJ*, 468, 462
 Haardt, F. & Madau, P. 1996, *ApJ*, 461, 20
 Haehnelt, M. G., Steinmetz, M., & Rauch, M. 1998, *ApJ*, 495, 647
 Hernquist, L., Katz, N., Weinberg, D. H., & Miralda-Escudé, J. 1996, *ApJ*, 457, L51
 Hogan, C. J. & Weymann, R. J. 1987, *MNRAS*, 225, 1P
 Hui, L. and Gnedin, N. Y., 1997, *MNRAS*, 292, 27
 Lowenthal, J. D., Hogan, C. J., Leach, R. W., Schmidt, G. D., & Foltz, C. B. 1990, *ApJ*, 357, 3
 Martínez-González, E., González-Serrano, J. I., Cayon, L., Sanz, J. L., & Martín-Mirónes, J. M. 1995, *A&A*, 303, 379
 McDonald, P., & Miralda-Escudé, J. 1999, *ApJ*, 519, 486
 Miralda-Escudé, J., & Ostriker, J. P. 1990, *ApJ*, 350, 1
 Miralda-Escudé, J., Cen, R., Ostriker, J. P., & Rauch, M. 1996, *ApJ*, 471, 582
 Mo, H. J. & Miralda-Escudé, J. 1996, *ApJ*, 469, 589
 Neufeld, D. A. 1990, *ApJ*, 350, 216
 Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: University Science Books)
 Péroux, C., McMahon, R. G., Storrie-Lombardi, L. J., & Irwin, M. J. 2003, *MNRAS*, 346, 1103
 Porciani, C., Magliocchetti, M., & Norberg, P. 2004, *MNRAS*, 355, 1010
 Prochaska, J. X., & Wolfe, A. M. 1997, *ApJ*, 474, 140
 Ryu, D., Ostriker, J. P., Kang, H., & Cen, R. 1993, *ApJ*, 414, 1
 Scott, J., Bechtold, J., Dobrzycki, A., & Kulkarni, V. P. 2000, *ApJS*, 130, 67
 Steidel, C. C., Bowen, D. V., Blades, J. C., & Dickenson, M. 1995, *ApJ*, 440, L45
 Stenflo, J. O. 1980, *A&A*, 84, 68
 Theuns, T., Leonard, A., Efstathiou, G., Pearce, F. R., & Thomas, P. A. 1998, *MNRAS*, 301, 478
 Zanstra, H. 1949, *Bull. Astron. Inst. Netherlands*, 11, 1
 Zhang, Y., Anninos, P., & Norman, M. L. 1995, *ApJ*, 453, L57
 Zheng, Z., & Miralda-Escudé, J. 2002a, *ApJ*, 568, L71
 Zheng, Z., & Miralda-Escudé, J. 2002b, *ApJ*, 578, 33